

Problem Solving

Consider a homogeneous isotropic molecular bar of circular cross-section, 20 nm in diameter, 2 micrometers long, with a Young's modulus of 1 GPa.

What is its bending spring constant?

$$\begin{array}{l}
 E = 10^9 \frac{N}{m^2} \\
 L = 2 \cdot 10^{-6} m \\
 D = 20 \cdot 10^{-9} m \\
 R = 10 \cdot 10^{-9} m \\
 \hline
 \kappa_b = ?
 \end{array}
 \quad
 \begin{array}{l}
 \kappa_b = \frac{4\pi}{3} \cdot \frac{E \cdot R^4}{L^3} = \\
 = \frac{4\pi}{3} \cdot \frac{10^9 \frac{N}{m^2} \cdot (10 \cdot 10^{-9} m)^4}{(2 \cdot 10^{-6} m)^3} = \\
 = \frac{\pi}{6} \cdot \frac{10^{-5} N}{m} = \\
 = 0.52 \cdot 10^{-5} N/m = 0.52 \cdot 10^{-2} pN/nm
 \end{array}$$

Consider a homogeneous isotropic molecular bar of circular cross-section, 20 nm in diameter, 2 micrometers long, with a Young's modulus of 1 GPa.

What is its tension spring constant?

$$\begin{array}{l}
 E = 10^9 \frac{N}{m^2} \\
 L = 2 \cdot 10^{-6} m \\
 D = 20 \cdot 10^{-9} m \\
 R = 10 \cdot 10^{-9} m \\
 \hline
 \kappa = ?
 \end{array}
 \quad
 \begin{array}{l}
 \kappa = \frac{E \cdot A}{L} = \\
 = \frac{E \cdot \pi R^2}{L} = \\
 = \frac{10^9 \frac{N}{m^2} \cdot \pi \cdot (10 \cdot 10^{-9} m)^2}{2 \cdot 10^{-6} m} = \\
 = \frac{\pi}{2} \cdot \frac{10^{-1} N}{m} = 157 pN/nm
 \end{array}$$

Consider a homogeneous isotropic molecular bar of circular cross-section, 20 nm in diameter, 2 micrometers long, with a Young's modulus of 1 GPa.

What is the persistence length at 27 °C?

$$\begin{array}{l}
 L = 2 \cdot 10^{-6} m \\
 E = 10^9 \frac{N}{m^2} \\
 D = 20 \cdot 10^{-9} m \\
 t = 27 \text{ } ^\circ\text{C} \\
 R = 10 \cdot 10^{-9} m \\
 T = 300 K \\
 \hline
 L_p = ?
 \end{array}
 \quad
 \begin{array}{l}
 L_p = \frac{E \cdot I}{k_B \cdot T} \quad I = \frac{\pi R^4}{4} \\
 L_p = \frac{10^9 \frac{N}{m^2} \cdot \frac{\pi}{4} \cdot (10 \cdot 10^{-9} m)^4}{1.38 \cdot 10^{-23} \frac{Nm}{K} \cdot 300 K} = \\
 = \frac{\pi \cdot 10^{-2} m}{1.38 \cdot 12} = 1.9 \cdot 10^{-3} m
 \end{array}$$

What is the smallest resolvable distance in a microscope that uses 520 nm light for imaging and has an aperture angle of 140 °?

$\lambda = 520 \text{ nm}$		$\delta = 0.61 \cdot \frac{\lambda}{n \cdot \sin \omega} =$
$2\omega = 140^\circ$		$= 0.61 \cdot \frac{520 \text{ nm}}{1 \cdot \sin 70^\circ} =$
$\omega = 70^\circ$		$= 0.61 \cdot \frac{520 \text{ nm}}{0.94} = 337 \text{ nm}$
$\delta = ?$		

How does the minimum distance that can be resolved by a microscope change if an immersion oil with a refractive index $n = 1.5$ is used?

$n = 1.5$		$\delta_n = 0.61 \cdot \frac{\lambda}{n \cdot \sin \omega} = \frac{\delta}{n} = \frac{\delta}{1.5}$
δ without immersion oil		
δ_n with immersion oil		
$\delta_n \quad ? \quad \delta$		

What is the minimum resolvable distance in a microscope that uses 420 nm light for imaging and has a numerical aperture of 1.4?

$\lambda = 420 \text{ nm}$		$\delta = 0.61 \cdot \frac{\lambda}{NA} =$
$NA = 1.4$		$= 0.61 \cdot \frac{420 \text{ nm}}{1.4} =$
$\delta = ?$		$= 0.61 \cdot 300 \text{ nm} = 183 \text{ nm}$